



**Patient Information:** Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Gender:  M  F  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_ Phone: \_\_\_\_\_ Alt. Phone: \_\_\_\_\_  
 Email: \_\_\_\_\_ Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ Care Giver: \_\_\_\_\_ Allergies:  No Known  See Attached Charts

**Prescriber Information:** Name: \_\_\_\_\_ Address: \_\_\_\_\_ City: \_\_\_\_\_  
 State: \_\_\_\_\_ Zip: \_\_\_\_\_ NPI: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Office Contact: \_\_\_\_\_

**Statement of Medical Necessity (Please Attach All Medical Documentation)**

**Date of Diagnosis:** \_\_\_\_\_ **ICD-10:** \_\_\_\_\_ **Other:** \_\_\_\_\_ **TB Test:**  Positive  Negative Date: \_\_\_\_\_  
**LFT:** ALT: \_\_\_\_\_ AST: \_\_\_\_\_ Date: \_\_\_\_\_ **Assessment:**  Moderate  Mod to Severe  Severe \_\_\_\_\_ % BSA Affected  
 Scalp  Face  Chest  Arms  Hands  Nails  Back  Groin  Buttocks  Legs  Other \_\_\_\_\_ **ISGA or EASI** \_\_\_\_\_

**Patient also taking methotrexate?**  Yes  No  
**Serious or active infection present?**  Yes  No  
**Hep B ruled out or treatment started?**  Yes  No  
**Does patient have latex allergy?**  Yes  No

**Injection Training:**  Pharmacist to Provide Training  
 Patient Trained in MD Office  
 Manufacturer Nurse Support  
 To Be Administered By a HCP

**Product Delivery:**  Patient's Home  
 Physician's Office  Pharmacy to Coordinate

Prior Failed Treatments	Drug Name	Length of Treatment
<input type="checkbox"/> 5-ASA		
<input type="checkbox"/> Biologics		
<input type="checkbox"/> Corticosteroids		
<input type="checkbox"/> Immunosuppressants		
<input type="checkbox"/> Methotrexate		
<input type="checkbox"/> NSAIDS		
<input type="checkbox"/> Surgery		
<input type="checkbox"/> Topical/Oral Antibiotics		
<input type="checkbox"/> UVA <input type="checkbox"/> UVB		
<input type="checkbox"/> Other		

**Prescription Information**

Medication	Dosage & Strength	Direction	Qty	Ref
<input type="checkbox"/> Dupixent	<b>Pediatric Atopic Dermatitis</b> <input type="checkbox"/> 300mg/2ml Prefilled Syringe <input type="checkbox"/> 200mg/1.14ml Prefilled Syringe <input type="checkbox"/> 300mg/2ml Prefilled Pen <i>(only for 12 years and older)</i>	<b>Induction Dose:</b> <input type="checkbox"/> ≥ 60 kg: Inject 600mg SC (two 300mg injections) <input type="checkbox"/> 30 to <60 kg: Inject 400mg SC (two 200mg injections) <input type="checkbox"/> 15 to <30 kg: Inject 600mg SC (two 300mg injections) <b>Maintenance Dose:</b> <input type="checkbox"/> ≥ 60 kg: Inject 300mg SC every other week <input type="checkbox"/> 30 to <60 kg: Inject 200mg SC every other week <input type="checkbox"/> 15 to <30 kg: Inject 300mg SC every 4 weeks	2	0
<input type="checkbox"/> Humira	<b>Hidradenitis Suppurativa</b> <input type="checkbox"/> Adolescent Hidradenitis Suppurativa 80mg/0.8ml and 40mg/0.4ml Starter Pack <input type="checkbox"/> Adolescent Hidradenitis Suppurativa 40mg/0.4ml Starter Package <input type="checkbox"/> Hidradenitis Suppurativa 80mg/0.8ml Starter pack <input type="checkbox"/> Hidradenitis Suppurativa 40mg/0.4ml Starter pack <input type="checkbox"/> 40mg/0.4ml Pen 40mg/0.4ml Prefilled Syringe	<b>Induction Dose:</b> <input type="checkbox"/> <b>Adolescents 12 years and older 66 lbs to &lt;132 lbs:</b> Inject 80mg SC on day 1, then 40mg SC on day 8 and every otherweek thereafter <input type="checkbox"/> <b>Adolescents 12 years and older &gt;132 lbs:</b> Inject two 80mgpens SC on day 1, then one 80mg pen SC on day 15 <input type="checkbox"/> <b>Adolescents 12 years and older &gt;132 lbs:</b> Inject one 80mgpen SC on day 1, then 80mg pen SC on day 2, then one 80mg pen SC on day 15 <b>Maintenance Dose:</b> <input type="checkbox"/> <b>Adolescents 12 years and older 66 lbs to &lt;132 lbs:</b> Inject40mg every other week <input type="checkbox"/> <b>Adolescents 12 years and older &gt;132 lbs:</b> Inject 40mg onday 29 then inject 40mg every week	3 4 3 6	0 0 0 0
<input type="checkbox"/> Humira	<b>Juvenile Idiopathic Arthritis + Pediatric Uveitis</b> <input type="checkbox"/> 10mg/0.1ml Prefilled Syringe <input type="checkbox"/> 20mg/0.2ml Prefilled Syringe <input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe	<input type="checkbox"/> <b>22 lbs to &lt;33 lbs:</b> Inject 10mg SC every other week <input type="checkbox"/> <b>33 lbs to &lt;66 lbs:</b> Inject 20mg SC every other week other week <input type="checkbox"/> <b>≥66 lbs:</b> Inject 40mg SC every	2	
<input type="checkbox"/> Cosentyx	<b>&lt;110lb</b> <input type="checkbox"/> 75mg Syringe <b>&gt;110lb</b> <input type="checkbox"/> 150mg Prefilled Syringe <input type="checkbox"/> 150mg Sensor Ready Pen	<b>Adolescents 6 years old and up &lt;110lb:</b> <input type="checkbox"/> <b>Induction Dose:</b> One 75mg injection under the skin at week 0, 1, 2, 3, 4. <input type="checkbox"/> <b>Maintenance Dose:</b> One 75mg injection under the skin every 4 weeks <input type="checkbox"/> <b>&gt;110lb: Induction Dose:</b> One 150mg injection under the skin at week 0, 1, 2, 3, 4. <input type="checkbox"/> <b>Maintenace Dose:</b> One 150mg injection under the skin every 4 weeks		

**Prescriber Signature** (I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs)

Signature: \_\_\_\_\_ Signature \_\_\_\_\_ Date: \_\_\_\_\_

**Substitution Permitted** **Dispense as Written**

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.



**Patient Information:** Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Gender:  M  F Last 4 of SSN: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_ Phone: \_\_\_\_\_ Alt. Phone: \_\_\_\_\_  
 Email: \_\_\_\_\_ Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ Care Giver: \_\_\_\_\_ Allergies:  No Known  See Attached Charts

**Prescriber Information:** Name: \_\_\_\_\_ Address: \_\_\_\_\_ City: \_\_\_\_\_  
 State: \_\_\_\_\_ Zip: \_\_\_\_\_ NPI: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Office Contact: \_\_\_\_\_

**Statement of Medical Necessity (Please Attach All Medical Documentation)**

**Date of Diagnosis:** \_\_\_\_\_ **ICD-10:** \_\_\_\_\_ **Other:** \_\_\_\_\_ **TB Test:**  Positive  Negative Date: \_\_\_\_\_  
**LFT:** ALT: \_\_\_\_\_ AST: \_\_\_\_\_ Date: \_\_\_\_\_ **Assessment:**  Moderate  Mod to Severe  Severe \_\_\_\_\_ % BSA Affected  
 Scalp  Face  Chest  Arms  Hands  Nails  Back  Groin  Buttocks  Legs  Other \_\_\_\_\_ **ISGA or EASI** \_\_\_\_\_

**Patient also taking methotrexate?**  Yes  No  
**Serious or active infection present?**  Yes  No  
**Hep B ruled out or treatment started?**  Yes  No  
**Does patient have latex allergy?**  Yes  No

**Injection Training:**  Pharmacist to Provide Training  
 Patient Trained in MD Office  
 Manufacturer Nurse Support  
 To Be Administered By a HCP

**Product Delivery:**  Patient's Home  
 Physician's Office  Pharmacy to Coordinate

Prior Failed Treatments	Drug Name	Length of Treatment
<input type="checkbox"/> 5-ASA		
<input type="checkbox"/> Biologics		
<input type="checkbox"/> Corticosteroids		
<input type="checkbox"/> Immunosuppressants		
<input type="checkbox"/> Methotrexate		
<input type="checkbox"/> NSAIDS		
<input type="checkbox"/> Surgery		
<input type="checkbox"/> Topical/Oral Antibiotics		
<input type="checkbox"/> UVA <input type="checkbox"/> UVB		
<input type="checkbox"/> Other		

**Prescription Information**

Medication	Dosage & Strength	Directions	Qty	Ref
<input type="checkbox"/> Humira	<b>Pediatric Crohn's Disease</b> <input type="checkbox"/> Pediatric Crohn's Starter Pack: 80mg/0.8ml, 40mg/0.4ml <input type="checkbox"/> Pediatric Crohn's Starter Pack: 80mg/0.8ml <input type="checkbox"/> 20mg/0.2ml Prefilled Syringe <input type="checkbox"/> 40mg/0.4ml Pen <input type="checkbox"/> 40mg/0.4ml Prefilled Syringe	<b>Induction Dose:</b> <input type="checkbox"/> <b>37 lbs to &lt;88 lbs:</b> Inject one 80mg pen SC on day 1, then one 40mg pen SC on day 15 <input type="checkbox"/> <b>&gt;88 lbs:</b> Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 <input type="checkbox"/> <b>&gt;88 lbs:</b> Inject one 80mg pen SC on day 1, then 80mg pen SC on day 2, then one 80mg pen SC on day 15	2	0
		<b>Maintenance Dose:</b> <input type="checkbox"/> <b>37 lbs to &lt;88 lbs:</b> Inject 20mg SC every other week <input type="checkbox"/> <b>&gt;88 lbs:</b> Inject 40mg SC every other week	3	0
			2	
<input type="checkbox"/> Stelara	<b>Pediatric Plaque Psoriasis</b> <input type="checkbox"/> 45mg/0.5ml Prefilled Syringe <input type="checkbox"/> 45mg/0.5ml Single-Dose Vial <input type="checkbox"/> 90mg/ml Prefilled Syringe	<b>Induction Dose:</b> <input type="checkbox"/> <b>&lt;60kg:</b> Inject 0.75mg/kg SC at week 0 <input type="checkbox"/> <b>60-100kg:</b> Inject 45mg SC at week 0 <input type="checkbox"/> <b>&gt;100kg:</b> Inject 90mg SC at week 0	1	0
		<b>Maintenance:</b> <input type="checkbox"/> <b>&lt;60kg:</b> Inject 0.75mg/kg at week 4, then every 12 weeks thereafter <input type="checkbox"/> <b>60-100kg:</b> Inject 45mg SC at week 4, then every 12 weeks thereafter <input type="checkbox"/> <b>&gt;100kg:</b> Inject 90mg SC at week 4, then every 12 weeks thereafter	1	0
			1	0
<input type="checkbox"/> Taltz	<b>Pediatric Plaque Psoriasis</b> <input type="checkbox"/> 80mg/ml Single-Dose Prefilled Autoinjector <input type="checkbox"/> 80mg/ml Single-Dose Prefilled Syringe  20mg and 40mg doses for patients weighing <=50kg (110lb) must be prepared and administered by a qualified healthcare professional.	<b>Induction Dose:</b> <input type="checkbox"/> <b>&gt;50kg:</b> Inject 160mg SC (two 80mg injections) at week 0 <input type="checkbox"/> <b>25 to 50kg:</b> Inject 80mg SC at week 0 <input type="checkbox"/> <b>&lt;25kg:</b> Inject 40mg SC at week 0	2	0
		<b>Maintenance:</b> <input type="checkbox"/> <b>&gt;50kg:</b> Inject 80mg SC at week 4 and every 4 weeks thereafter <input type="checkbox"/> <b>25-50kg:</b> Inject 40mg SC at week 4 and every 4 weeks thereafter <input type="checkbox"/> <b>&lt;25kg:</b> Inject 20mg SC at week 4 and every 4 weeks thereafter	1	
<input type="checkbox"/>				

**Prescriber Signature** (I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs)

Signature: \_\_\_\_\_ Signature: \_\_\_\_\_ Date: \_\_\_\_\_

**Substitution Permitted** **Dispense as Written**

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.

**Confidentiality Notice:** This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.