



Patient Information: Name: _____ DOB: _____ Gender: M F Last 4 of SSN: _____
 Address: _____ City: _____ State: _____ Zip: _____ Phone: _____ Alt. Phone: _____
 Email: _____ Ht: _____ Wt: _____ Care Giver: _____ Allergies: No Known See Attached Charts

Prescriber Information: Name: _____ Address: _____ City: _____
 State: _____ Zip: _____ NPI: _____ Phone: _____ Fax: _____ Office Contact: _____

Statement of Medical Necessity (Please Attach All Medical Documentation)

Date of Diagnosis: _____ **ICD-10:** _____ **Other:** _____ **TB Test:** Positive Negative **Date:** _____

Patient also taking Methotrexate? Yes No
 Serious or active infection present? Yes No
 Hep B ruled out or treatment started? Yes No
 Does patient have latex allergy? Yes No

LFT: AST: _____ ALT: _____ **Date:** _____

Injection Training: Pharmacist to Provide Training
 Patient Trained in MD Office Manufacturer Nurse Support

Product Delivery: Patient's Home Physician's Office
 Pharmacy to Coordinate

Prior Failed Treatments	Drug Name and Length of Treatment
<input type="radio"/> Azulfidine	
<input type="radio"/> Biologics	
<input type="radio"/> Calcipotriene	
<input type="radio"/> Celebrex	
<input type="radio"/> Corticosteroids	
<input type="radio"/> Indocin	
<input type="radio"/> Methotrexate	
<input type="radio"/> Other	

Prescription Information:

Medication	Dosage & Strength	Directions	Qty	Ref
<input type="radio"/> Otezla	<input type="radio"/> Two-Week Starter Pack (Titration) <input type="radio"/> 28-Day Starter Pack (Titration) <input type="radio"/> 30mg Tablets	<input type="radio"/> Starter Pack: Take one tab in the morning on Day 1, then take one tab in the morning and one tab in the evening as directed per pack	1	
		<input type="radio"/> Maintenance: Take one 30mg Tablet by mouth twice daily <i>For patients with severe renal impairment take one 30mg tablet once daily and skip afternoon doses in starter pack.</i>	60	
<input type="radio"/> Rasuvo	<input type="radio"/>	<input type="radio"/>		
<input type="radio"/> Rinvoq	<input type="radio"/> 15mg Extended Release Tablets	<input type="radio"/> Take one 15mg Tablet once a day	30	
<input type="radio"/> Simponi	<input type="radio"/> 50mg/0.5ml Smartject Autoinjector <input type="radio"/> 50mg/0.5ml Prefilled Syringe	<input type="radio"/> Inject 50mg SC once a month	1	
<input type="radio"/> Stelara (for PsA)	<input type="radio"/> 45mg/0.5ml Prefilled Syringe <input type="radio"/> 45mg/0.5ml Vial <input type="radio"/> 90mg/1ml Prefilled Syringe <input type="radio"/> Stelara self-injection: <i>Healthcare provider certifies that patient has been trained and is eligible for self-injection</i>	<input type="radio"/> Induction Dose: Inject 45mg SC on day one	1	
		<input type="radio"/> Maintenance: Inject 45mg SC on day 29, and every 12 wks thereafter	1	
		<input type="radio"/> PsA with Coexistent Moderate to Severe Plaque Psoriasis (>220lb) <input type="radio"/> Induction Dose: Inject 90mg SC on day 1	1	
		<input type="radio"/> Maintenance: Inject 90mg SC on day 29, and every 12 wks thereafter	1	
<input type="radio"/> Taltz	<input type="radio"/> 80mg/ml Single-Dose Prefilled Autoinjector <input type="radio"/> 80mg/ml Single-Dose Prefilled Syringe	Ankylosing Spondylitis <input type="radio"/> Induction Dose: Inject 160mg SC (two 80mg injections) at week 0	2	0
		<input type="radio"/> Maintenance: Inject 80mg SC every 4 weeks	1	
		Non-Radiographic Axial Spondyloarthritis <input type="radio"/> Inject 80mg SC every 4 weeks	1	
		Psoriatic Arthritis: <input type="radio"/> Induction Dose: Inject 160mg SC (two 80mg injections) at week 0	2	0
		<input type="radio"/> Maintenance: Inject 80mg SC every 4 weeks	1	
		PsA with Coexistent Moderate to Severe Plaque Psoriasis <input type="radio"/> Weeks 0-2: Inject 160mg SC (two 80mg injections) at week 0, then inject 80mg at week 2	3	0
		<input type="radio"/> Weeks 4-10: Inject 80mg SC at week 4 and every 2 weeks thereafter through week 10	2	1
		<input type="radio"/> Weeks 12 & Onward: Inject 80mg SC at week 12 and every 4 weeks thereafter	1	
Tremfya	<input type="radio"/> 100mg/ml Prefilled Syringe <input type="radio"/> 100mg/ml One-Press Patient Controlled Injector	<input type="radio"/> Induction Dose: Inject 100mg SC at weeks 0 and 4	2	0
		<input type="radio"/> Maintenance: Inject 100mg SC every 8 weeks thereafter	1	
Xeljanz Xeljanz XR	<input type="radio"/> 5mg Tablet <input type="radio"/> 11mg Tablet	<input type="radio"/> Take one 5mg tablet by mouth twice a day	60	
		<input type="radio"/> Take on 11mg tablet by mouth once a day *For patients with moderate renal or hepatic impairment take one 5mg tablet once daily*	30	

Prescriber Signature (I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs)

Signature: _____ Signature: _____ Date: _____

Substitution Permitted **Dispense as Written**

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment.

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.